



# N.C. Department of Agriculture and Consumer Services



N.C. Forest Service



Hemlock conservation area at Hanging Rock State Park, Stokes County, NC.

## Annual Legislative Report on Hemlock Restoration

Oct. 1, 2023

Steve Troxler, Commissioner  
Scott Bissette, Assistant Commissioner  
David Lane, State Forester

Pursuant to G.S. 106-927, the North Carolina Forest Service, a division of the North Carolina Department of Agriculture and Consumer Services, respectfully submits this annual report on hemlock restoration in North Carolina for fiscal year 2023.

## Contents

STATUTORY REQUIREMENT .....	3
§ 106-927. Annual report on hemlock restoration .....	3
IMPORTANCE OF HEMLOCK FORESTS TO NORTH CAROLINA.....	3
AN OVERVIEW OF HEMLOCK WOOLLY ADELGID INFESTATION AND CONTROL MEASURES IN NORTH CAROLINA .....	5
HEMLOCK RESTORATION INITIATIVE GOALS AND ACCOMPLISHMENTS: 2014 - PRESENT .....	9
HEMLOCK RESTORATION INITIATIVE ACCOMPLISHMENTS: FY 2023.....	9
CONTACT INFORMATION.....	13
Appendix A .....	16
Appendix B.....	17
Appendix C.....	18

**\*Cover: Hemlock conservation area at Hanging Rock State Park, Stokes County, NC.**

## STATUTORY REQUIREMENT

### § 106-927. Annual report on hemlock restoration


Beginning Oct. 1, 2022, and no later than Oct. 1 of each year, the Commissioner shall submit a written report on hemlock restoration in North Carolina to the Joint Legislative Oversight Committee on Agriculture and Natural and Economic Resources and the Fiscal Research Division. The report shall include the following with respect to each hemlock restoration initiative funded during the previous fiscal year:

- (1) Identification of goals and outcomes for the initiative.
- (2) A description of the measures used, or data collected to evaluate the efficiency and effectiveness of the initiative in reaching its desired goals and outcomes.
- (3) The performance of each initiative with respect to the identified goals and outcomes.

## IMPORTANCE OF HEMLOCK FORESTS TO NORTH CAROLINA

Hemlock trees have a wide range extending south from New Brunswick and Nova Scotia in Canada to Alabama along the Appalachian Mountains and as far west as Minnesota, with isolated populations existing east and west of the normal range. In North Carolina, hemlocks are a significant part of the 18.8 million acres of forestland. It occurs throughout the Appalachian Mountain region, descending into the foothills and areas in the Piedmont. A disjunct occurrence is present in the Hemlock Bluffs Nature Preserve in Cary (the reason for the preserves formation) approximately 100 miles east of the edge of the normal range. Two species of hemlock occur in North Carolina; the eastern hemlock (*Tsuga canadensis*) and the Carolina hemlock (*Tsuga caroliniana*). Eastern hemlock dominates the wide range described above. In contrast, Carolina hemlock is a Southern Appalachian endemic, restricted to western North

Carolina, with smaller populations occurring in Virginia and Georgia. North Carolina is home to roughly 80% of the naturally occurring Carolina hemlock, a unique position for our state.



Hemlocks are foundational species in the forest ecosystems of our state and beyond. They're a long-lived tree species capable of obtaining a large size. It is not unusual for hemlock trees to live for hundreds of years, with the oldest known individuals exceeding 800 years. One of the largest eastern hemlocks on record obtained a diameter of 76 inches and a height of 175 feet. Hemlocks have long been referred to as "the redwood of the east."

A foundational species is requisite to the survival of other plants and animals in these forests. In the case of hemlock, the shade provided by the overstory canopy is responsible for maintaining a microclimate in the understory conducive to the survival of many other plants and animals. In addition, hemlock is shade tolerant and can maintain a full crown, even under the cover of other trees including other hemlocks. These characteristics make it a vital source of winter cover for various wildlife species such as ruffed grouse, white-tailed deer and eastern cottontail rabbit. American black bear commonly use eastern hemlock cavities for winter denning. It also provides critical shading for mountain streams by maintaining temperatures adequate to support many aquatic organisms including several species of salamanders and native brook trout. Indeed, hemlock forests are essential to the survival of more than 90 different bird species, 120 vertebrates and numerous plant communities, some of which are listed as threatened or endangered on the state or federal level. Included here are the blackpoll warbler, eastern hellbender and piratebush.



Figure 2 left to right:  
Piratebush photo by Will Cook.  
Blackpoll warbler photo by Simon Boivin/Maccaulay Library.  
Eastern hellbender photo by N.C. Wildlife Resources Commission.

Hemlock forests also have a direct significance on many human activities. Many think of forested areas being used for campsites and recreation areas. The loss of hemlock in these areas impacts people by creating hazard trees that must be removed at a substantial cost or pose a direct threat to public safety as they fall. The visual impact of these dead trees also influences the large tourism industry within the hemlock range. Additionally, it has been reported that dead hemlock trees falling into the

waterways in which they tend to grow have contributed to destructive flooding events in the mountains in recent years.




Figure 3: Carolina hemlock at high elevations. NC is home to approximately 80% of the world's population of this species.

## **AN OVERVIEW OF HEMLOCK WOOLLY ADELGID INFESTATION AND CONTROL MEASURES IN NORTH CAROLINA**

Hemlock woolly adelgid (HWA) is an aphid like insect native to Asia and the west coast of North America where it was first identified in 1928. It is a native pest in its natural range and has an adapted suite of natural predators that keep populations in check under normal conditions. It was first detected in Richmond, VA in the early 1950's. The original infestation was determined to have originated from Honshu Island, Japan.

Hemlock woolly adelgid has a complex life cycle consisting of two phases known as the progrediens (spring) and sistens (fall/overwinter) generations. The sistens generation is active and feeds during the winter. This is the time when most pesticide treatments occur. Although HWA has both males and females in its native range, only females exist in eastern North America. In their native range, where sexual reproduction occurs, males feed on tiger-tail spruce. This portion of the life cycle has not been shown to occur in eastern North America as our native red spruce does not seem to support male HWA. In eastern North America, HWA has an abbreviated life cycle where females are parthenogenic, or able to reproduce asexually.



Figure 4: Early reinfestation of hemlock woolly adelgid following treatment, Little Fork HCA. Inset: A hemlock woolly adelgid showing its feeding tube. Photo by Kelly Oten, NCSU.



Figure 5: Healthy hemlocks in Wilkes County, NC. These trees were treated during the 2018-2019 fiscal year.

As eggs hatch following the progrediens generation, the early nymphs, known as crawlers, travel to the base of the needles, settle and enter a period of dormancy. Around October, they put on their woolly covering, which aids in protection and shelters their eggs. They feed, mature and lay their eggs in the early spring. Adults from these eggs begin the progrediens generation again. Timing within this life cycle is critical for treatment operations (Appendix A).

In 1995, HWA was detected in Stokes and Surry counties. The invasion expanded steadily following entry to the point that most of the hemlock range in North Carolina was infested by 2007. That same year, widespread mortality was noticed in the earlier infested areas (Appendix B).

Following initial detection, control methods centered around trees in urban locations or areas heavily traveled by people such as campgrounds. Much of the original response consisted of outreach and provision of insecticides and technical advice to those desiring to treat hemlocks. This was done by N.C. Forest Service (NCFS) forest health professionals and other selected NCFS personnel.

The NCFS began treatments on state lands in 2007, with the first treatments taking place at Dupont State Recreational Forest (DSRF). The following year, NCFS facilitated the treatment of trees on South Mountains State Park and in 2009 assisted the N.C. Wildlife Resources Commission (NCWRC) with treatments on several game land properties. These efforts resulted in approximately 5,000 hemlocks being treated directly through NCFS efforts.

In 2011, the NCFS was transferred from the N.C. Department of Environment and Natural Resources to the N.C. Department of Agriculture and Consumer Services (NCDA&CS). Following that transfer, the plight of hemlocks in our state became a focal objective for the Agricultural Commissioner Steve Troxler, culminating in the formation of the Hemlock Restoration Initiative (HRI) as a signature program in 2014. This program is operated under the nonprofit WNC Communities and has been funded through NCDA&CS, federal grants obtained by NCFS and state appropriations. A large portion of the control of HWA and hemlock restoration was transferred to HRI following the formation of the program.

Hemlock restoration continues to be a cooperative effort between NCDA&CS and WNC Communities. Within NCDA&CS, NCFS and the Beneficial Insects Lab under the Plant Industry Division are the main cooperators. Numerous other state and federal agencies and private partners are involved including the N.C. Division of Parks and Recreation, the NCWRC, the US Department of Agriculture Forest Service (USFS) and many others. The remainder of this report will focus on accomplishments related to the HRI culminating in a summary of accomplishments from the 2022-2023 fiscal year.

## HEMLOCK RESTORATION INITIATIVE GOALS AND ACCOMPLISHMENTS: 2014 – PRESENT

### *Hemlock Restoration Initiative Goals*

The Hemlock Restoration Initiative Goals and Measures of Accomplishment are:

- Identifying and establishing hemlock conservation areas (HCA) – *Number of HCAs established.*
- Educating landowners on how to economically treat and manage the hemlocks on their properties – *Number of outreach programs, social media posts and engagement and additional trees treated on private lands.*
- Increasing the number of trees being treated on public lands – *Number of trees treated and cumulative inches of diameter (breast height) treated.*
- Implementing Integrated Pest Management (IPM) and long-term biological control of HWA – *Number of sites and counties in which HWA predators recovered. Number of release sites for HWA predators.*
- Advancing the development of other control strategies and restoration techniques including the search for HWA resistant trees and the growing conditions that hemlocks like best – *Number of impact plots established or revisited. Pounds of hemlock cones collected. Number of hemlock seedlings produced.*

### *Overall Accomplishments*

Throughout the life of this initiative, work has been completed related to the objectives mentioned above. A summary of these objectives is as follows:

- 185 Hemlock Conservation Areas established on state, private and federal lands.
  - 28 separate state properties are represented in this list.
- All hemlock conservation areas have received Phase I treatments (chemical). Areas scheduled for return treatments are being evaluated for Phase II treatments.
  - Phase II treatments are based on the presence of beneficial predators of HWA and consist of partial chemical treatments to promote the growth of beneficial insect populations.
- Approximately 120,000 hemlocks have been chemically treated.
- 246 educational outreach events were conducted.
- 913 individual inquiries from landowners and the public serviced.
- 7,447 trees are known to have been treated as a result of this information sharing.
- 1,956 volunteers have been recruited resulting in 6,153 service hours.
- 22,248 predatory *Laricobius* beetles released for biocontrol across North Carolina.
- More than 7,370 *Laricobius* beetles recovered from 130 locations, many not documented release sites.

- *Laricobius* beetles documented in 17 of the 27 counties within the hemlock range.
- Five *Laricobius* insectary sites established, three on state lands.
- Collection of beetles and support for NCDA&CS Plant Industry Division's Beneficial Insects Lab and University of Tennessee's Lindsay Young Beneficial Insect Lab.
- Raising of three separate species of predatory beetles within North Carolina.
- More than 900 pounds of hemlock cones collected and processed for seed production.
- Establishment of a Carolina hemlock seed collection area.
- Successful rearing of both Carolina and eastern hemlock seedlings at Linville River Nursery for public sale.
- More than 90,000 hemlock seedlings were produced for sale.



Figure 6: NCFS Forest Health Specialist Brian Heath working with a B.R.I.D.G.E. crew member during 2023 treatment operations.

## **HEMLOCK RESTORATION INITIATIVE ACCOMPLISHMENTS: 2022-2023 FISCAL YEAR**

Throughout the 2022-2023 fiscal year, North Carolina's HRI continued to work toward achieving its goals. The three major entities involved, the NCFS, NCDA&CS Plant Industry Beneficial Insects Lab and WNC Communities HRI, worked together and in concert with partners to achieve the following:

- 20,707 hemlocks treated chemically.
  - Largest known population of Carolina hemlock treated.
- 30 hemlock conservation areas with Phase II treatments.
- 16 new hemlock conservation areas established.
- 11 new impact plots were established to measure the effects of treatments over time.
- 26 educational and outreach programs organized and presented.
- 51 partner/stakeholder meetings organized and/or attended.
- 126 social media posts generated.
- 12,593 social media engagements from the public.
- 158 hemlock inquiries serviced.
- 1,447 additional hemlocks were treated by private individuals as a result of outreach activities.
- Temporary beneficial insect technicians funded allowing additional collections.
- Percival I36-VLC8 growth chamber utilized for the rearing of *Laricobius* and other HWA predators.
- Survey and collection of *Sasajiscymnus tsugae* (released 2002-2015) indicating a viable population of this predatory insect.
- 46 days of monitoring/assessment for biocontrol program.
- *Laricobius* documented in 18 sites over nine counties indicating widespread establishment.
- 1,173 *Laricobius* adults/eggs released (992 *L. ozakensis* -potentially the most effective species).
- 52,515 hemlock seedling cells sown (39,150 eastern hemlock/13,365 Carolina hemlock).
- Collection of 297 lbs. of eastern hemlock and 8 lbs. of Carolina hemlock cones for seedling production.
- Planting of eastern and Carolina hemlock for seed production at Morganton Forestry Center.



Figure 7: HRI staff conducting a treatment workshop for private landowners and state parks personnel. Hanging Rock State Park, 2023.



Figure 8: Cooperative planting of hemlocks for seed production, Morganton Forestry Center.




Figure 9: NCDA&CS Beneficial Insects Lab research study site.



Figure 10: HRI treatment crew in action.



Figure 11: Sampling for beneficial insects at the Bald Knob Hemlock Conservation Area.



Figure 12: Carolina hemlock seedling planted at Morganton Forestry Center.

## CONTACT INFORMATION

For further information, please contact:

Jim Slye  
N.C. Forest Service - Forest Health Branch Head  
N.C. Department of Agriculture and Consumer Services  
[james.slye@ncagr.gov](mailto:james.slye@ncagr.gov)  
919-857-4858  
1616 Mail Service Center, Raleigh, NC 27699-1600


Amanda Poole  
N.C. Forest Service - Administrative Services Division Director  
N.C. Department of Agriculture and Consumer Services  
[amanda.poole@ncagr.gov](mailto:amanda.poole@ncagr.gov)  
919-857-4853  
1616 Mail Service Center, Raleigh, NC 27699-1600

## Appendix A

# Hemlock Woolly Adelgid (*Adelges tsugae*) Life Cycle



## Appendix B



## Appendix C

